



<https://doi.org/10.5800/GT-2017-8-3-0279>

Proceedings of the Second Russia–China International Meeting on the
Central Asian Orogenic Belt (September 6–12, 2017, Irkutsk, Russia)

THE EARLY-MIDDLE PALEOZOIC VOLCANISM AND GEODYNAMIC EVOLUTION OF THE HERLEN MASSIF, CENTRAL PART OF THE CAOB: CONSTRAINS FROM GEOCHEMISTRY, U-Pb GEOCHRONOLOGY, LU-Hf AND Rb-Sr ISOTOPES OF VOLCANIC ROCKS

Ts. Narantsetseg¹, D. Orolmaa¹, Chao Yuan², Tao Wang³,
Lei Guo³, Ying Tong³, Xinyu Wang², T. Oyunchimeg¹,
O. Enkh-Orshikh¹, P. Delgerzaya¹

¹*Institute of Paleontology and Geology, Mongolian Academy of Sciences,
Ulaanbaatar, Mongolia*

²*State Key Laboratory of Isotope Geochemistry, Guangzhou Institute of Geochemistry,
Chinese Academy of Sciences, Guangzhou 510640, China*

³*Institute of Geology, Chinese Academy of Geological Sciences, Beijing 100037, China*

For citation: Narantsetseg Ts., Orolmaa D., Yuan C., Wang T., Guo L., Tong Y., Wang X., Oyunchimeg T., Enkh-Orshikh O., Delgerzaya P., 2017. The Early-Middle Paleozoic volcanism and geodynamic evolution of the Herlen massif, central part of the CAOB: Constrains from geochemistry, U-Pb geochronology, Lu-Hf and Rb-Sr isotopes of volcanic rocks. *Geodynamics & Tectonophysics* 8 (3), 521–523. doi:10.5800/GT-2017-8-3-0279.

Mongolia lies in the central part of the Central Asian Orogenic Belt [Mossakovsky *et al.*, 1994; Zorin, 1999; Jahn, 2004; Khain *et al.*, 2003; Badarch *et al.*, 2002; Windley *et al.*, 2007; Zhang *et al.*, 2008], or Altaids [Şengör *et al.*, 1993; Şengör, Natal'in, 1996; Wilhem *et al.*, 2012], which is fringed by the Siberian craton in the north and by the Tarim and Sino-Korean Cratons in the south. According to the recent tectonic subdivision, the territory of Mongolia is subdivided into Northern and Southern domains which are separated by the so called Mid Mongolian Tectonic Line [Tomurtogoo,

2012]. The Herlen Massif is one of the important tectonic units of the South Mongolian domain in the Argun-Idermeg super terrane extending through the territories of Russia and China [Parfenov *et al.*, 2009; Tomurtogoo, 2014b]. The Herlen massif, also known as Herlen superterrane [Tomurtogoo, 2012] or Idermeg terrane [Tomurtogoo, 2014a] is composed of Ereendavaa, Undur-Khaan, Idermeg and Gobian Altay-Baruun Urt terranes converged at the end of the Cambrian-beginning of the Ordovician [Badarch *et al.*, 2002; Tomurtogoo, 2014b].

The Ereendavaa terrane is situated in extreme northeast and contains metamorphic rocks of Paleoproterozoic and Mesoproterozoic Khaychin-gol and Ereendavaa formations overlying by Late Neoproterozoic-Early Cambrian formations [Marinov et al., 1973; Blagonravov et al., 1990; Dorjnamjaa, Bat-Ireedui, 1991; Byamba, 1991; Dorjnamjaa et al., 2011; Tomurtogoo, 2012]. Lower Paleozoic granitoids and Late Ordovician to Late Carboniferous marine sedimentary and terrigenous-volcanic rocks widely distribute in this terrane [Tomurtogoo, 2012].

In this paper, we present new geochronological, geochemical and isotopic data for the Late Neoproterozoic-Early Cambrian volcanic rocks from Tsarigiin gol area, southern part of Ereendavaa terrane in order to examine their formation time, petrogenesis and tectonic environment.

The Tsarigiin gol area is located in the south of the Ereendavaa terrane, bordered by the Undurkhaan terrane in the southeast. Middle to Upper Neoproterozoic Zamttolgoi metamorphic complex, mainly composed of gneiss, various schists, quartzite, and minor amphibolite, are the oldest stratum in this area [Narantsetseg et al., 2015]. The Zamttolgoi metamorphic complex is overlain by the thin basal conglomerate and schistose sandstone, followed by volcanic successions erupted at Lower Paleozoic [Kalimulin et al., 1968; Chongradi et al., 1985], or Neoproterozoic-Early Cambrian [Makhbadar, Delgertsogt, 1990] without precise dating and detailed geochemical data.

The 2000–3500 m thick volcanic strata is well exposed around the Undurkhaan Mountain in the Tsarigiin gol area. Three sections were identified. Section I is a rhyolitic composition, and consists of only felsic volcanic rocks. Section II is characterized by intermediate to felsic volcanic rocks which were intensively subjected to green schist facies metamorphism. Section III has a bimodal distribution and is composed of mainly felsic and minor mafic volcanic rocks.

A subset of 18 samples was collected from I–III sections to obtain mineralogical and geochemical characteristics of volcanic rocks Tsarigiin gol area in the Ereendavaa terrane. A total of 3 rhyolite samples were dated by LA-ICPMS, two of them were analyzed for Hf isotopic compositions. Also, Sr and Nd isotopic compositions of one mafic and three intermediate volcanic rock samples were used for this study.

Zircons from rhyolite samples previously believed to be Neoproterozoic-Early Cambrian, are euhedral-subhedral in shape and show fine-scale oscillatory

growth zoning and high Th/U ratios ranging between 0.47 and 1.24, indicating a magmatic origin. LA-ICPMS zircon U-Pb dating indicate that the volcanic activity in the southern part of Ereendavaa terrane took place in two stages: Late Ordovician (~462–455 Ma), and Early Devonian (~418 Ma) instead of Neoproterozoic-Early Cambrian or Lower Paleozoic. The Late Ordovician volcanic rocks are intermediate to felsic in composition, and characterized by an enrichment of the LILE and LREE and depletion of HFSE, accompanied by positive $\epsilon_{\text{Hf}}(t)$ and $\epsilon_{\text{Nd}}(t)$ values and slightly negative $\epsilon_{\text{Hf}}(t)$ values. Early Devonian volcanic rocks display a typical bimodal distribution in composition with dominant rhyolite and minor trachybasalt and basaltic trachyandesite. Basalts are medium K-calc-alkaline, whereas felsic end member is high K-calc-alkaline and display the geochemical characteristics of A2-type granites and characterized by mostly negative $\epsilon_{\text{Hf}}(t)$ and $\epsilon_{\text{Nd}}(t)$ values. The geochemical similarities of Late Ordovician and Early Devonian volcanic rocks are implying that magmas were generated at subduction zone which is consistent with the geology of area. But, spatial and temporal distribution of volcanic rocks and their different composition together with some geochemical characteristics are indicating that these rocks were originated in different tectonic setting. The Late Ordovician volcanic rocks probably formed in continental arc setting developed in active continental margin, by extensive fractional crystallization of basaltic magmas with minor involvement of Mesoproterozoic continental crust, whereas Early Devonian bimodal volcanic rocks occur in post collisional extensional rift setting. Basalts were derived from high degrees partial melting of lithospheric mantle and the felsic rocks may derived by partial melting of crustal material with involvement of juvenile mantle materials.

Combined our new data, a Neoproterozoic to Paleozoic tectonic evolution model of southern part of Ereendavaa terrane can be set. Neoproterozoic to Late Ordovician, active continental margin setting developed in the southern part of Ereendavaa terrane and northwestward subduction (present coordinates) of Undurkhaan oceanic plate beneath the Ereendavaa continental terrane took place with medium to high-K calc-alkaline volcanic eruption in the Tsarigiin gol area. In the Late Silurian, Ereendavaa and Idermeg terranes collided, and the post collisional Early Devonian bimodal volcanic association with A2-type felsic end-members distributed in the southern margin of Ereendavaa terrane.

REFERENCES

- Badarch G., Cunningham D.W., Windley B.F., 2002. A new terrane subdivision for Mongolia: implications for the Phanerozoic crustal growth of Central Asia. *Journal of Asian Earth Sciences* 21 (1), 87–110. [https://doi.org/10.1016/S1367-9120\(02\)00017-2](https://doi.org/10.1016/S1367-9120(02)00017-2).

- Blagonravov B.I., Goldenberg V.I., Blagonravova L.A. et al., 1990. Report of 1:200000 geological mapping, State Geological Fund of Mineral Resources and Petroleum Authority of Mongolia, No 1757, Ulaanbaatar (in Russian).
- Byamba J., 1991. Tectonic Evolution of Mongolia in Late Proterozoic-Early Paleozoic, D.Sc. Thesis. Geological Institute, Russian Academy of Sciences, Moscow, 350 p. (in Russian).
- Chongradi E., Peregi J., Dienish E. et al., 1985. Report of 1:50000 geological mapping, State Geological Fund of Mineral Resources and Petroleum Authority of Mongolia, No 3797, Ulaanbaatar (in Russian).
- Dorjnamjaa D., Bat-Ireedui Ya., 1991. Precambrian of Mongolia. Ulaanbaatar, 182 p. (in Russian).
- Dorjnamjaa D., Byamba J., Enkhbaatar B., 2011. Precambrian of Mongolia. In: Geology and mineral resources of Mongolia, vol. 1, Stratigraphy, p. 52–99.
- Jahn B.M., 2004. The Central Asian Orogenic Belt evolution and growth of the continental crust in the Phanerozoic. In: J. Malpas, C.J.N. Fletcher, J.R. Ali, J.C. Aichison (Eds.), Aspects of the Tectonic Evolution of China. Geological Society, London, Special Publications, vol. 226, p. 73–100. <https://doi.org/10.1144/GSL.SP.2004.226.01.05>.
- Kalimulin S.M., Dyrante M.B. et al., 1968. Report of 1:200000 geological mapping, State Geological Fund of Mineral Resources and Petroleum Authority of Mongolia, No1758, Ulaanbaatar (in Russian).
- Khain E.V., Bibikova E.V., Salnikova E.B., Kröner A., Gibsher A.S., Didenko A.N., Degtyarev K.E., Fedotova A.A., 2003. The Palaeo-Asian ocean in the Neoproterozoic and Early Paleozoic: New geochronologic data and palaeotectonic reconstructions. *Precambrian Research* 122 (1–4), 329–358. [https://doi.org/10.1016/S0301-9268\(02\)00218-8](https://doi.org/10.1016/S0301-9268(02)00218-8).
- Makhbadar Ts., Delgertsogt B., 1990. Explanatory note for the geology map of Central and Eastern Mongolia, scale 1:500000. Ulaanbaatar (in Mongolian).
- Marinov N.A., Zonenshain L.P., Blagonravov V.A. (Eds.), 1973. Geology of the Mongolian People's Republic, vol. 1, Stratigraphy. Nedra, Moscow, 782 p. (in Russian).
- Mossakovsky A.A., Ruzhentsev S.V., Samygin S.G., Kheraskova T.N., 1994. Central Asian fold belt: geodynamic evolution and formation history. *Geotectonics* 27 (6), 445–474.
- Narantsetseg Ts., Chao Y., Tomurhuu D., Delgerzaya P., Enkh-Orshikh O., 2015. Geochronology, geochemistry of metamorphic rocks from Herlen area. *Explorer* 53, 69–88 (in Mongolian).
- Parfenov L.M., Badarch G., Berzin N.A., Khanchuk A.I., Kuzmin M.I., Nokleberg W.J., Prokopiev A.V., Ogasawara M., Yan H., 2009. Summary of Northeast Asia geodynamics and tectonics. In: Stephan Mueller Special Publication Series, vol. 4, p. 11–33. <http://www.stephan-mueller-spec-publ-ser.net/4/11/2009/>.
- Sengör A.M.C., Natal'in B.A., 1996. Paleotectonics of Asia: fragments of synthesis. In: A. Yin, T.M. Harrison (Eds.), The tectonic evolution of Asia. Cambridge University Press, Cambridge, p. 486–640.
- Sengör A.M.C., Natal'in B.A., Burtman V.S., 1993. Evolution of the Altaid tectonic collage and Palaeozoic crustal growth in Eurasia. *Nature* 364 (6435), 299–306. <https://doi.org/10.1038/364299a0>.
- Tomurtogoo O., 2012. Tectonic subdivision of orogenic belt of Mongolia. *Transactions of Institute of Geology and Mineral Resources of MAS, Geology* 21, 5–25 (in Mongolian).
- Tomurtogoo O., 2014a. The structure and genesis of old massifs of Mongolia. *Geology* 28, 23–33 (in Mongolian).
- Tomurtogoo O., 2014b. Tectonics of Mongolia. In: Tectonics of Northern, Central and Eastern Asia, explanatory note to the Tectonic map of Northern Central Eastern Asia and adjacent areas at scale 1:2500000, p. 110–126.
- Wilhem C., Windley B.F., Stampfli G.M., 2012. The Altaids of Central Asia: a tectonic and evolutionary innovative review. *Earth-Science Reviews* 113 (3–4), 303–341. <https://doi.org/10.1016/j.earscirev.2012.04.001>.
- Windley B.F., Alexeiev D., Xiao W., Kröner A., Badarch G., 2007. Tectonic models for accretion of the Central Asian Orogenic Belt. *Journal of the Geological Society* 164 (1), 31–47. <https://doi.org/10.1144/0016-76492006-022>.
- Zhang X., Zhang H., Tang Y., Wilde S.A., Hu Z., 2008. Geochemistry of Permian bimodal volcanic rocks from central Inner Mongolia, North China: Implication for tectonic setting and Phanerozoic continental growth in Central Asian Orogenic Belt. *Chemical Geology* 249 (3–4), 262–281. <https://doi.org/10.1016/j.chemgeo.2008.01.005>.
- Zorin Yu.A., 1999. Geodynamics of the western part of the Mongolia-Okhotsk collisional belt, Trans-Baikal region (Russia) and Mongolia. *Tectonophysics* 306 (1), 33–56. [https://doi.org/10.1016/S0040-1951\(99\)00042-6](https://doi.org/10.1016/S0040-1951(99)00042-6).